Diagnosis Report



PATIENT ID AGE GENDER

1521 Female

INTERVIEW SOURCE

INTERVIEW DATE

Youth

05/04/2018

Current Diagnosis

Depressive Disorders

Major Depressive Disorder, moderate, single episode (F32.1)

Depressed Mood, Present

Anhedonia, Present

Patient reported trouble falling asleep or staying asleep More than half the days the past two weeks. Problem began April 2017

Psychomotor Agitation in Depressive Disorder, Present

Decreased Self-Esteem, Present

Concentration Disturbance, Present

Indecision, Present

Impairment in functioning due to depression, Present

Persistent Depressive Disorder (Dysthymia) with persistent major depressive episode PRESENT F34.1

Depressed Mood, Present

Anhedonia, Present

Patient reported trouble falling asleep or staying asleep More than half the days the past two weeks. Problem began April 2017

Decreased Self-Esteem, Present

Concentration Disturbance, Present

Indecision, Present

Hopeless, Present

Impairment in functioning due to depression, Present

Generalized Anxiety Disorder

Generalized Anxiety Disorder – Present (F41.1)

Excessive worries more days than not, Present

Excessive worries across breadth of domains, Present ()

Worry associated with defined symptom(s), Present (i feel restless, i have difficulty concentrating, i have difficulty falling asleep, i have difficulty staying asleep,)

Difficulty controlling worries, Present

Impairment in functioning due to worries, Present

Clinically significant distress due to worries, Present

Worrying has lasted at least 6 months, Present

Sleep Problems

SLEEP PROBLEMS, Present

Patient reported trouble falling asleep or staying asleep More than half the days the past two weeks. Problem began April 2017

Suicidality

Suicidality, Present

Symptom	Description	C-CASA Code	Patient Comments
Self-Injurious Behavior without suicidal intent	Self-injurious behavior with no intent to die. The behavior is intended purely for other reasons.	7	

Past Diagnosis

Suicidality

Suicidality, Past

Symptom	Description	C-CASA Code	Patient Comments
Past Suicidal ideation:	Past thoughts of wanting to	4	

	2017)		
icidal ideation: tive, method	Past thoughts of suicide and thought of at least one method,	4	

Copyright © 2023 KSADS-COMP Inc. 12/13/2023 11:35 AM